

Matrix-Product Ansatz for Excited States of Fractional Quantum Hall Systems

Zheng-Yuan WANG and Masaaki NAKAMURA

Department of Physics, Tokyo Institute of Technology, O-Okayama, Meguro-ku, Tokyo 152-8551, Japan

E-mail: wang@stat.phys.titech.ac.jp, masaaki@stat.phys.titech.ac.jp

(Received January 24, 2013)

Recently, it was discussed that the $\nu = 1/3$ fractional quantum Hall state can be expressed by a one-dimensional lattice model with an exact matrix-product ground state, when toroidal boundary conditions are assumed for a narrow strip [Phys. Rev. Lett. **109** (2012) 016401]. In this article, we discuss how the excitation spectra of this system are calculated analytically based on the matrix-product formalism. We introduce variational wave functions for various momenta and optimize them. The obtained spectra of the charge neutral excitations show the magneto-roton behavior.

KEYWORDS: fractional quantum Hall effect, thin-torus limit, matrix product state, exactly solvable model, excited states, magneto-roton

1. Introduction

In recent years, there have been theoretical efforts to study the fractional quantum Hall (FQH) states around the thin-torus (or Tao-Thouless, TT) limit, where the original 2D system in a magnetic field with toroidal boundary conditions is reduced to 1D lattice models with short-range interactions [1–4]. This approach is justified because the FQH states in this anisotropic system are known to be adiabatically connected to the isotropic cases [2]. From this point of view, the authors with Bergholtz have shown that the $\nu = 1/3$ Laughlin state [5] can be expressed by a 1D model with an exact matrix-product (MP) ground state [6]. In this article, we give the details to obtain the excitation spectra of this model based on the MP wave function with variational parameters which partially mentioned in Ref. [6], and generalize this calculation to arbitrarily momenta and obtain roton behavior of the excitation spectra [7].

2. Fractional quantum Hall states on torus geometries

We consider a model of N_e interacting electrons in a magnetic field B on a torus boundary conditions with circumferences L_1 and L_2 for x and y -directions, respectively. For simplicity, we normalize x, y, L_1, L_2 by magnetic length $l_B = \sqrt{\hbar/eB}$. When the torus is pierced by N_s magnetic flux quanta, boundary conditions require the relation $L_1 L_2 = 2\pi N_s$. In Landau gauge ($\mathbf{A} = By\hat{x}$), a complete basis of N_s degenerate single-particle states in the lowest Landau level can be chosen as

$$\psi_i(x, y) = \frac{1}{\sqrt{\pi^{1/2} L_1}} \sum_{j \in \mathbb{Z}} \exp \left[i \left(\frac{2\pi}{L_1} i + L_2 j \right) x \right] \exp \left[-\frac{1}{2} \left(y + \frac{2\pi}{L_1} i + L_2 j \right)^2 \right], \quad (1)$$

where the guiding center of the Gaussian function in y -direction is given by the quantized momentum in x -direction with $i = 0, 1, \dots, N_s - 1$. In this basis, any translationally-invariant 2D Hamiltonian with two-body interactions assumes the following 1D lattice model,

$$\mathcal{H} = \sum_{|m| < k \leq N_s/2} \hat{V}_{km}, \quad \hat{V}_{km} \equiv \sum_i V_{km} c_{i+m}^\dagger c_{i+k}^\dagger c_{i+k+m} c_i, \quad (2)$$

where the matrix-element V_{km} specifies the amplitude for a pair hopping process where two particles separated $k + m$ sites hop m steps to opposite directions (note that m can be negative). The $m = 0$ terms are the electrostatic repulsions, the matrix elements are real and antisymmetric functions for interchange of k and m . This model conserves the center of mass coordinate defined by $K = \sum_{i=1}^{N_s} i n_i \pmod{N_s}$ with $n \equiv c_i^\dagger c_i$, which is identified as the total momentum in x -direction [see Eq. (1)].

For small L_1 the matrix elements V_{km} are simplified considerably: Since one finds that for generic interaction, the $m \neq 0$ terms are exponentially suppressed as

$$V_{km} \sim V_{k0} \exp(-2\pi^2 m^2 / L_1^2). \quad (3)$$

Therefore, in the TT limit $L_1 \rightarrow 0$, $m \neq 0$ terms in the original Hamiltonian (2) can be neglected, so that the Hamiltonian can be rewritten as $\mathcal{H}_{\text{TT}} = \sum_i \sum_j V_{k0} n_{i+j} n_j$. Then the ground state is obtained at filling factor $\nu = 1/3$ is gapped, and three-fold degenerate charge density wave state with three sites periodicity, $|\Psi_0\rangle = |010|010|010 \dots\rangle$. A system of thin but finite torus can be considered by taking the leading hopping terms \hat{V}_{km} ($m \neq 0$) into account for the TT state $|\Psi_0\rangle$.

3. Matrix-product ground state for $\nu = 1/3$ Laughlin state

The exactly solvable model for $\nu = 1/3$ states [6] can be obtained by truncating the Hamiltonian (2) up to the third nearest interactions ($k + |m| \leq 3$) as an approximation in the vicinity of the TT limit. When we further assume $L_2 \rightarrow \infty$ limit, the matrix elements of the pseudo potential for $\nu = 1/3$ Laughlin state [8] satisfies the conditions $V_{21}^2 = V_{10} V_{30}$ and $V_{20} > 0$. Then the Hamiltonian is rewritten as $\mathcal{H}_{1/3} = \sum_i [Q_i^\dagger Q_i + P_i^\dagger P_i]$, where $Q_i = \alpha_0 c_{i+1} c_{i+2} + \alpha_1 c_i c_{i+3}$ and $P_i = \beta_0 c_i c_{i+2}$ with $\alpha_0, \alpha_1, \beta_0 \in \mathbf{R}$. This Hamiltonian apparently has positive expectation values $\langle \mathcal{H}_{1/3} \rangle \geq 0$, and the following wave function as the exact ground state

$$|\Psi\rangle = \prod_i (1 + t c_{i+1}^\dagger c_{i+2}^\dagger c_{i+3} c_i) |\Psi_0\rangle, \quad t \equiv -\alpha_1 / \alpha_0, \quad (4)$$

since $Q_i |\Psi_{1/3}\rangle = P_i |\Psi_{1/3}\rangle = 0, \forall i$ is satisfied. This is three-fold degenerate but can be shown to be unique ground state for each center-of-mass sectors.

The wave function (4) can be written in a MP form $|\Psi_{1/3}\rangle = \text{tr}[g_1 g_2 \dots g_{N_e}]$ [9, 10] where the matrix g_i is identified in the following way. First, we introduce the $S = 1$ spin representation for the three-sites unit cell as $|010\rangle \rightarrow |o\rangle$, $|001\rangle \rightarrow |+\rangle$ and $|100\rangle \rightarrow |-\rangle$. One finds that the possible configurations of the nearest two spins are $|oo\rangle + t|+-\rangle + |o+\rangle + |-o\rangle + |-\rangle$, so that if we express the relations in the following 3×3 matrix

$$g_i = \begin{bmatrix} f_{--} |-\rangle_i & f_{-o} |o\rangle_i & f_{-+} |+\rangle_i \\ f_{o-} |-\rangle_i & f_{oo} |o\rangle_i & f_{o+} |+\rangle_i \\ f_{+-} |-\rangle_i & f_{+o} |o\rangle_i & f_{++} |+\rangle_i \end{bmatrix}, \quad (5)$$

its non-vanishing elements are identified as $f_{oo} = f_{o+} = f_{-o} = f_{-+} = 1$ and $f_{+-} = t$. Next, we reduce the obtained 3×3 matrix to a 2×2 form by changing the base from $(t|-\rangle, |o\rangle, |+\rangle)$

to $(t|-\rangle + |o\rangle, |+\rangle)$ as

$$g_i \equiv \begin{bmatrix} |o\rangle_i & |+\rangle_i \\ t|-\rangle_i & 0 \end{bmatrix}. \quad (6)$$

The matrix (6) has a 4×4 transfer matrix $G \equiv \bar{g}_j \otimes g_j$ with only two eigenvalues $\lambda_{\pm} = (1 \pm \sqrt{4t^2 + 1})/2$. The argument for the Hamiltonian with an exact MP wave function can also be extended to general $\nu = 1/q$ Laughlin states [11].

4. Variational wave functions for excited states

We derive excitation spectra at $\nu = 1/3$ by a variational method. In the TT limit, the charge neutral excited states include one 01-type and one 0-type domain walls that carry fractional charges $e^* = e/3$ and $e^* = -e/3$, respectively. These states $|\Psi_0^{\Delta K=1}\rangle = |010|0|01|010|010 \dots\rangle$, $|\Psi_0^{\Delta K=2}\rangle = |010|0|010|01|010 \dots\rangle$, \dots , are classified by the center of mass coordinate relative to the ground state which is related to the momentum of the x -direction. Now we consider the variational wave functions that consists of configurations generated from $|\Psi_0^{\Delta K}\rangle$ by applying the Hamiltonian $\mathcal{H}_{1/3}$. For $\Delta K = 1$, there are following exchange processes, $|010|001|010|010 \dots\rangle \rightarrow |010|001|001|100 \dots\rangle \rightarrow |010|000|110|100 \dots\rangle \rightarrow |010|001|001|100 \dots\rangle$. Since the interactions in $\mathcal{H}_{1/3}$ are limited within the third neighbors, there is no exchange process across the defects. We can rewrite these processes in the $S = 1$ spin variables by inserting vacancies appropriately (between the two 1's) [12, 13] as

$$\begin{aligned} |010|0|010|010|010 \dots\rangle &\rightarrow |010|0|010|001|100 \dots\rangle \rightarrow \\ |010|0|010|010|010 \dots\rangle &\rightarrow |010|0|010|001|100 \dots\rangle. \end{aligned} \quad (7)$$

We redefine the site indices so that the defect comes to the edge, and introduce an ansatz with variational parameters u and v as $|\Psi^{\Delta K=1}\rangle = \text{tr}[g'_1 g_2 g_3 \dots g_{N_e-1} g''_{N_e}]$, where

$$g'_1 \equiv \begin{bmatrix} u|o\rangle_1 & 0 \\ t|-\rangle_1 & v|o\rangle_1 \end{bmatrix}, \quad g''_{N_e} \equiv \begin{bmatrix} |o\rangle_{N_e} & 0 \\ t|-\rangle_{N_e} & 0 \end{bmatrix}. \quad (8)$$

For the case $\Delta K \geq 2$, the ansatz should include three variational parameters u , v and w as $|\Psi^{\Delta K}\rangle = \text{tr}[g_1 g_2 \dots g_{K-1} g'''_K g_{K+2} \dots g_{N_e-1} g''_{N_e}]$ with

$$g'''_i \equiv \begin{bmatrix} |oo\rangle_{i,i+1} & u|o+\rangle_{i,i+1} + v|+o\rangle_{i,i+1} \\ tu|o-\rangle_{i,i+1} + tv|o-\rangle_{i,i+1} & tw|-\rangle_{i,i+1} \end{bmatrix}. \quad (9)$$

When we calculate the compressibility, we need to shrink the system size as $N_s \rightarrow N_s - 1$ by removing 0 from the root state. Then only one 01-type domain wall ($|\dots 010|010|010 \dots\rangle$) should be considered, so that the ansatz becomes $|\Psi^-\rangle = \text{tr}[g'''_1 g_3 g_4 \dots g_{N_e-1}]$.

These variational wave functions for $N_e \rightarrow \infty$ can be optimized analytically by using the undetermined coefficient method. Fig. 1 (a) shows the excitation spectra obtained by the MP ansatz for the infinite system as functions of L_1 . These well agree with the results obtained from the infinite systems by exact diagonalization (ED). The deviation in the large L_1 region is due to the finite size effect which appears through λ_-/λ_+ . We also get a finite value of $E(N_s - 1)$ for $N_e \rightarrow \infty$, reflecting that the Laughlin state is incompressible $\kappa^{-1} = \lim_{N_s \rightarrow \infty} (N_s/4\pi l_B^2)[E(N_s - 1) + E(N_s + 1) - 2E_s(N_s)] \rightarrow \infty$ where $E(N_s + 1) = E_s(N_s) = 0$. Fig. 1 (b) shows the neutral excitation spectra obtained by the MP ansatz for $\Delta K = 1-6$. The $\Delta K = 2$ state is always the lowest, and the excitation energy converges to a constant value when ΔK is increased as shown in the inset. This captures a character of the magneto-roton behavior in the FQH state [7]. These results are also consistent with a recent analysis based on spherical coordinate and the Jack polynomials [14].




Fig. 1. (a) Comparison of excitation spectra obtained by the MP ansatz for infinite systems and those by exact diagonalization (ED) for a finite system $N_s = 27$. (b) Neutral excitation spectra for different momenta $\Delta K = 1-6$ derived from the MP ansatz for an infinite system. The inset shows the excitation spectra as a function of ΔK at $L_1 = 8$ which show a magneto-roton minimum.

5. Conclusion

We have studied the excitation spectra of the $\nu = 1/3$ FQH states considering toroidal boundary conditions in the vicinity of the Tao-Thouless limit. In this region, the original 2D system can be mapped onto a 1D lattice model with short-range interactions which has the exact MP ground state. We have introduced “ansatz” for the excited state as the MP state with defects and variational parameters, and derived the excitation spectra analytically. The excitation spectrum with a fractional charge reflects the incompressibility of fractional quantum Hall states. The charge neutral excitations behave like the roton mode.

6. Acknowledgment

We thank Emil J. Bergholtz for the related collaboration in Ref. [6]. Z.-Y. W. and M. N. acknowledge support from the Global Center of Excellence Program “Nanoscience and Quantum Physics” of the Tokyo Institute of Technology. M. N. also acknowledges support from Grant-in-Aid No.23540362 by MEXT.

References

- [1] R. Tao and D. J. Thouless: Phys. Rev. B **28** (1983) 1142.
- [2] E. H. Rezayi and F. D. M. Haldane: Phys. Rev. B **50** (1994) 17199.
- [3] E. J. Bergholtz and A. Karlhede: Phys. Rev. Lett. **94** (2005) 026802.
- [4] A. Seidel, H. Fu, D. -H. Lee, J. M. Leinaas and J. Moore: Phys. Rev. Lett. **95** (2005) 266405.
- [5] R. B. Laughlin: Phys. Rev. Lett. **50** (1983) 1395.
- [6] M. Nakamura, Z. -Y. Wang and E. J. Bergholtz: Phys. Rev. Lett. **109** (2012) 016401.
- [7] S. M. Girvin, A. H. MacDonald and P. M. Platzman: Phys. Rev. Lett. **54** (1985) 581; Phys. Rev. B **33** (1986) 2481.
- [8] S. A. Trugman and S. A. Kivelson: Phys. Rev. B **31** (1985) 5280.
- [9] M. Fannes, B. Nachtergale and R. F. Werner: Europhys. Lett. **10** (1989) 633; Commun. Math. Phys. **144** (1992) 443.
- [10] A. Klümper, A. Schadschneider and J. Zittartz: Z. Phys. B **87** (1992) 281; Europhys. Lett. **24** (1993) 293.
- [11] Z. -Y. Wang and M. Nakamura: arXiv:1206.3071v2.
- [12] M. Nakamura, Z. -Y. Wang and E. J. Bergholtz: J. Phys.: Conf. Ser. **302**, 012020 (2011).
- [13] Z. -Y. Wang, S. Takayoshi and M. Nakamura: Phys. Rev. B **86** (2012) 155104.
- [14] B. Yang, Z.-X. Hu, Z. Papić and F. D. M. Haldane: Phys. Rev. Lett. **108** (2012) 256807.